



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Draft Genome Sequence of an *Enterococcus faecalis* Strain Isolated from a Neonatal Blood Sepsis Patient

Citation for published version:

Kropp, KA, Lucid, A, Carroll, J, Belgrudov, V, Walsh, P, Kelly, B, Smith, C, Dickinson, P, O'Driscoll, A, Templeton, K, Ghazal, P & Sleator, RD 2014, 'Draft Genome Sequence of an *Enterococcus faecalis* Strain Isolated from a Neonatal Blood Sepsis Patient', *Genome announcements*, vol. 2, no. 5.
<https://doi.org/10.1128/genomeA.00907-14>

Digital Object Identifier (DOI):

[10.1128/genomeA.00907-14](https://doi.org/10.1128/genomeA.00907-14)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Genome announcements

Publisher Rights Statement:

Copyright © 2014 Kropp et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Draft Genome Sequence of an *Enterococcus faecalis* Strain Isolated from a Neonatal Blood Sepsis Patient

K. A. Kropp,^a A. Lucid,^a J. Carroll,^b V. Belgrudov,^c P. Walsh,^b B. Kelly,^b C. Smith,^d P. Dickinson,^d A. O'Driscoll,^c K. Templeton,^{d,e} P. Ghazal,^d R. D. Sleator^a

Department of Biological Sciences, Cork Institute of Technology, Bishopstown, Cork, Ireland^a; NSilico Lifescience Ltd., Cork, Ireland^b; Department of Computing, Cork Institute of Technology, Bishopstown, Cork, Ireland^c; Division of Pathway Medicine, University of Edinburgh, Edinburgh, Scotland, United Kingdom^d; Microbiological Diagnostic Unit, Royal Infirmary, University of Edinburgh, Edinburgh, Scotland, United Kingdom^e

Herein, we report the draft genome sequence of *Enterococcus faecalis* ED-NGS-1009, cultivated from a blood sample taken from a neonatal sepsis patient at the Royal Infirmary in Edinburgh, Scotland, United Kingdom.

Received 11 August 2014 Accepted 20 August 2014 Published 11 September 2014

Citation Kropp KA, Lucid A, Carroll J, Belgrudov V, Walsh P, Kelly B, Smith C, Dickinson P, O'Driscoll A, Templeton K, Ghazal P, Sleator RD. 2014. Draft genome sequence of an *Enterococcus faecalis* strain isolated from a neonatal blood sepsis patient. *Genome Announc.* 2(5):e00907-14. doi:10.1128/genomeA.00907-14.

Copyright © 2014 Kropp et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to R. D. Sleator, roy.sleator@cit.ie.

Enterococcus faecalis is a nosocomial, opportunistic pathogen showing resistance to antibiotics, specifically vancomycin (1–4). Preterm neonates are a highly susceptible patient group for bacterial infections (1, 5, 6) and rapid detection of blood sepsis and the causative pathogen are critical steps to enable proper treatment (7–9). Therefore, in the ClouDx-i project, we are aiming to extend our existing knowledge of currently circulating pathogenic strains linked with blood sepsis in neonates to inform the development of new molecular diagnostic assays. Herein, we present the draft genome sequence of an *Enterococcus faecalis* strain isolated from a preterm neonate at the Royal Infirmary in Edinburgh, in 2013. Positivity for blood sepsis and species identification were confirmed by classical microbiological techniques.

The isolate was grown overnight at 37°C on Luria broth (LB) agar, and genomic DNA was isolated using Qiagen genomic tips (Venlo, Limburg, Netherlands). Genomic DNA was fragmented (fragments 2 to 10 kb) using sonication and a non-size selected genome library was produced using the Nextera mate pair kit (Illumina, San Diego, CA). This library was then sequenced on an Illumina MiSeq using MiSeq Reagent kit version 3. Genomic sequence assembly, analysis, and automated reporting were carried out using Simplicity (10). This approach produced 2,954,539 total reads, resulting in an average 222-fold coverage. The average G+C content was 37.27%. Sequence assembly was achieved using a *de novo* assembly pipeline based on the SPAdes version 3.10 assembly tool with k-mers K21, K33, K55, K77, K99, and K127, resulting in a total of 79 contigs, of which 13 were >1,000 bp, representing 99.02% of the total sequence information with the largest contig being 2,285,020 bp in size. Postassembly processing was performed using SPAdes, and only scaffolds of >1,000 bp were considered when estimating genome size as 2,973,146 bp. We annotated the genome with Prokka (11) and used the identified 16S rRNA gene to confirm the species as *Enterococcus faecalis*. A scaffold of the genome was produced with Contiguator2 and we identified the closest related strain by BLASTing the scaffold, returning *E. faecalis* V583 and DENG1 as similar but not identical isolates, as

was evident through insertions and deletions in the genome. The genome was then screened using GLIMMER3 (12) identifying 2,852 open reading frames (ORFs). The predicted ORFs were compared to the UniProt-TrEMBL database (13) using BLASTp, mapping 1,650 ORFs. To identify potential virulence factors in the genome, it was compared to a local database built from the VFDB (14) and Victors databases with BLASTp, using a 75% amino-acid sequence identity cutoff while only considering alignments longer than 100 amino-acids. In all, 63 potential virulence factors were identified.

Samples were handled in accordance with local ethical approval by the ethics committees of the NHS Lothian SAHSC Bioresource and the NHS Research and Development Office, Project ID 2011/R/NE/01 and the HSS BioResource Request ID 13/ES/0126.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JPWN00000000](https://www.ncbi.nlm.nih.gov/nuclseq/JAWN00000000/). The version described in this paper is version JPWN01000000.

ACKNOWLEDGMENT

This work was supported by the ClouDx-i IAPP EU FP7 project, coordinated by R. D. Sleator.

REFERENCES

- Kaufman D, Fairchild KD. 2004. Clinical microbiology of bacterial and fungal sepsis in very-low-birth-weight infants. *Clin. Microbiol. Rev.* 17: 638–680. <http://dx.doi.org/10.1128/CMR.17.3.638-680.2004>.
- Hasman H, Saputra D, Sicheritz-Ponten T, Lund O, Svendsen CA, Frimodt-Møller N, Aarestrup FM. 30 October 2013. Rapid whole-genome sequencing for the detection and characterization of microorganisms directly from clinical samples. *J. Clin. Microbiol.* <http://dx.doi.org/10.1128/JCM.02452-13>.
- Fisher K, Phillips C. 2009. The ecology, epidemiology and virulence of *Enterococcus*. *Microbiology* 155:1749–1757. <http://dx.doi.org/10.1099/mic.0.026385-0>.
- Bizzarro MJ, Raskind C, Baltimore RS, Gallagher PG. 2005. Seventy-five years of neonatal sepsis at Yale: 1928–2003. *Pediatrics* 116:595–602. <http://dx.doi.org/10.1542/peds.2005-0552>.

5. Shah BA, Padbury JF. 2014. Neonatal sepsis: an old problem with new insights. *Virulence* 5:170–178. <http://dx.doi.org/10.4161/viru.26906>.
6. Ghazal P, Dickinson P, Smith CL. 2013. Early life response to infection. *Curr. Opin. Infect. Dis.* 26:213–218. <http://dx.doi.org/10.1097/QCO.0b013e32835fb8bf>.
7. Labib AZ, Mahmoud AB, Eissa N, El Gendy FM, Soliman MA, Aly AA. 2013. Early diagnosis of neonatal sepsis: a molecular approach and detection of diagnostic markers versus conventional blood culture. *Int. J. Microbiol. Res.* 4:77–85.
8. Mancini N, Carletti S, Ghidoli N, Cichero P, Burioni R, Clementi M. 2010. The era of molecular and other non-culture-based methods in diagnosis of sepsis. *Clin. Microbiol. Rev.* 23:235–251. <http://dx.doi.org/10.1128/CMR.00043-09>.
9. Sibley CD, Peirano G, Church DL. 2012. Molecular methods for pathogen and microbial community detection and characterization: current and potential application in diagnostic microbiology. *Infect. Genet. Evol.* 12:505–521. <http://dx.doi.org/10.1016/j.meegid.2012.01.011>.
10. Walsh P, Carroll J, Sleator RD. 2013. Accelerating *in silico* research with workflows: a lesson in simplicity. *Comput. Biol. Med.* 43:2028–2035. <http://dx.doi.org/10.1016/j.combiomed.2013.09.011>.
11. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
12. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res.* 27:4636–4641. <http://dx.doi.org/10.1093/nar/27.23.4636>.
13. Boeckmann B, Bairoch A, Apweiler R, Blatter M-C, Estreicher A, Gasteiger E, Martin MJ, Michoud K, O'Donovan C, Phan I, Pilboud S, Schneider M. 2003. The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Res.* 31:365–370. <http://dx.doi.org/10.1093/nar/gkg095>.
14. Chen L, Yang J, Yu J, Yao Z, Sun L, Shen Y, Jin Q. 2005. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids Res.* 33:D325–D328. <http://dx.doi.org/10.1093/nar/gki008>.